arXiv:1302.7185v2 [quant-ph] 10 Mar 2013

Generalized Fermat principle for classical and quantum systems

Tarek A. Elsayed[1, 2, *]
[1] Institute for Theoretical Physics, University of Heidelberg, Philosophenweg 19, 69120 Heidelberg, Germany
[2] Electronics Research Institute, Dokki, Giza 12622, Egypt

The analogy between dynamics and optics had a great influence on the development of the foundations of classical and quantum mechanics. We take this analogy one step further and propose a new variational principle in classical phase space and quantum Hilbert space that mimics Fermat principle in optics. The state of a many-body classical (quantum) system follows a trajectory in the phase (Hilbert) space that extremizes the time functional. The classical Hamilton's equations of motion and the quantum Schrodinger's equation are shown to lead necessarily to a stationary time functional. A plethora of other unconventional variational principles related to the geometry of the dynamical evolution, and inspired from Fermat principle are discussed in this paper.

It has been a hallmark in the history of physics in the past few centuries that ingenious and elegant reconsiderations of the relations between geometric concepts and physical phenomena lead to paradigm shifts in physics and our perception of the universe. The most notable examples are the theories of special and general relativity. Novel reconsiderations of geometric quantities in physics often lead to new variational principles which assign the natural evolution of physical systems with an extremum of some functional or a geodesic curve in some hyperspace.

The quest to find new variational principles in physics has not stopped till today [1]. The oldest of these variational principles is Fermat principle of least time which became a fundamental principle in geometric optics. The principle was introduced by Fermat, which he also called *the principle of natural economy* [2], and states that light rays travel in a general medium along the path that minimizes the time taken to travel between the initial and final destinations. The concept of natural economy inspired Maupertuis to introduce the principle of least action in analytical mechanics which later evolved through the work of Euler, Lagrange, Hamilton, and Jacobi to become a fundamental and universal concept in classical mechanics.

The mathematical formulation of Fermat principle states that the time functional $\mathcal{T}$ defined as

$$\mathcal{T} = \int_{s_i}^{s_f} \frac{ds}{\nu(s)} \tag{1}$$

is minimized along the trajectory of light rays[3]. In other words, if $\mathcal{T}$ is computed along all possible trajectories between an initial and final positions s_i, s_f where $\nu(s)$ is the speed of light and ds is the distance element along the trajectory, $\mathcal{T}$ will always be minimum along the actual path travelled by the light rays. The modern version of Fermat principle is written in terms the index of refraction $n(s) = \frac{c}{\nu(s)}$ where c is the speed of light in free space and states that the optical path length $\int_{s_i}^{s_f} n(s)ds$ is a minimum. In that sense, Fermat principle is the optical analog of Jacobi least action principle [4], which states that for a conservative classical system at energy E, with potential function between its constituent particles V, the action functional

$$I = \int_{s_i}^{s_f} \sqrt{E - V(s)}ds \tag{2}$$

computed in the configuration space along the physical path traversed by the system between the initial and final representative points is an extremum. The remarkable property of this action that distinguishes it from other variational principles in analytical mechanics, i.e., Hamilton's and Lagrange's variational principles, is that it represents a purely geometric quantity that is computed along different trajectories in the configuration space between fixed points without referring to any evolution in time. Therefore, equation 2 can be used to define a new Riemannian space whose metric $ds' = \sqrt{E - V(s)}ds$ where the natural evolution of the representative point of the system will be along geodesic curves.

In this work we elevate the status of the time functional (1) to a universal action in both classical and quantum generic many-body systems that yields an extremum along the physical path in a proper space $\mathbb{S}$ that fully characterizes the state of the system and its future evolution, once the Hamiltonian is known. A proper space for quantum systems is the projective Hilbert space $\mathbb{P}$, where wavefunctions are defined upto an overall phase factor, and for classical systems a proper space is the phase space consisting of all the generalized coordinates and momenta $\{q_i, p_i\}$. The principle informally states that among all possible trajectories between s_i and s_f only those which make $\mathcal{T}$ invariant under an infinitesimal variation in the path, i.e.,

$$\delta\mathcal{T} = 0 \tag{3}$$

are possible candidates for the dynamical evolution.

The expression for the speed of the representative point of the system $\nu(s)$ computed over $\mathbb{S}$ is obtained from the equations of motion of the system that will drive its evolution along the physical path (i.e., Schrodinger's equation in quantum systems and Hamilton's equations of motion in classical systems). We emphasize here that we are not on the mission of deriving these equations of motion from the time action, because we have to use them to find $\nu(s)$ in the first place, but rather to assert that they necessarily lead to a stationary time action. Despite being a purely geometric quantity, the time functional by definition can straightforwardly extract the timing information of the evolution once the path is determined. However, Fermat principle in a generic many-body system is in general defined for a time-independent system, i.e., the Hamiltonian does not contain time explicitly. Nevertheless, we use insights gained from this principle to define a new variational principle for quantum mechanics which is valid for a generic time-dependent Hamiltonian. Our study opens the door for introducing new variational principles in other classes of physical systems not considered in this paper (e.g., continuum mechanics and field theory) and hopefully shed more light on the role of time in physics.

—*Fermat Principle in Hilbert Space* Unlike the theory of general relativity, where geometric concepts are rooted in the bones of the theory, quantum mechanics did not enjoy such a solid connection with geometry in its very beginning. Later development of the geometric phase in quantum mechanics has encouraged many researchers to pay more attention to this issue. Anandan and Aharonov have initiated the investigations of the nature of the geometry of quantum evolution in the projective Hilbert space $\mathbb{P}$ through a series of papers in the late 80s [5–7]. They have shown [6] that the speed of quantum evolution in $\mathbb{P}$ is related to the energy uncertainty $\Delta E = \left(\langle H^2\rangle - \langle H\rangle^2\right)^{1/2}$ via

$$ds = 2\Delta E \; dt/\hbar \tag{4}$$

where ds is the infinitesimal distance in $\mathbb{P}$ given by the Fubini-Study (FS) metric $ds^2 = \langle\delta\psi|\delta\psi\rangle - |\langle\delta\psi|\psi\rangle|^2$. The trajectory traversed by a ray in $\mathbb{P}$ under unitary evolution is not a geodesic, i.e., $\delta\int ds \neq 0$. This can be immediately envisaged by considering a system composed of a single quantum spin 1/2. In this case, $\mathbb{P}$ is simply the Bloch sphere and the precession motion of the spin on Bloch sphere is not a geodesic. Several attempts[8, 9] tried to conceive new formulations where the quantum evolution is a geodesic flow.

The simple answer to this problem suggested by the present paper is to consider $\mathcal{T} = \int \frac{ds}{\Delta E}$ as a geodesic quantity, which is stationary along the physical path through $\mathbb{P}$, i.e., *Fermat Principle in Hilbert Space.* This is to be distinguished from the quantum brachistochrone problem[10], where the Hamiltonian that leads to optimal time evolution between an initial and final state is sought. The above proposition states that the unitary evolution generated by any Hamiltonian is optimal, with respect to all other possible trajectories connecting the initial and final states (Fig. 1.a).

If we parametrize the evolution along any path connecting $|\psi_i\rangle$ and $|\psi_f\rangle$ by some arbitrary parameter τ, we can write equation 1 as

$$\mathcal{T} = \int_{|\psi_i\rangle}^{|\psi_f\rangle} d\tau \frac{\langle\dot{\psi}|1-\hat{P}|\dot{\psi}\rangle^{\frac{1}{2}}}{\left(\langle\psi|H^2|\psi\rangle - \langle\psi|H|\psi\rangle^2\right)^{\frac{1}{2}}} \tag{5}$$

where $|\dot{\psi}\rangle = \frac{|\delta\psi\rangle}{\delta\tau}$ and $\hat{P}$ is the projection operator $|\psi\rangle\langle\psi|$. Taking the variational derivative of $\mathcal{T}$ with respect to $\langle\delta\psi|$ subject to the constraints of normalization and fixed initial and final states, we arrive at the Euler-Lagrange (EL) equation of the integrand L,

$$\frac{\delta L}{|\delta\psi\rangle} - \frac{d}{d\tau}\frac{\delta L}{|\delta\dot{\psi}\rangle} = 0 \tag{6}$$

After taking the inner product with $|\psi\rangle$ and making use of the constraint condition $\langle\psi|\psi\rangle = 1$ we arrive finally at

$$\langle\dot{\psi}|\psi\rangle^2 = -\frac{\langle\dot{\psi}|\dot{\psi}\rangle - |\langle\psi|\dot{\psi}\rangle|^2}{\langle\psi|H^2|\psi\rangle - \langle\psi|H|\psi\rangle^2}\langle\psi|H|\psi\rangle^2 \tag{7}$$

We can see straightforwardly that the Schrodinger's equation $|\dot{\psi}\rangle = \pm iH|\psi\rangle$ is the only linear evolution for $|\psi(t)\rangle$ that will satisfy this condition. The fact that both the forward and backward time evolutions exist simultaneously on the energy shell for the same variational problem, i.e., between the same initial and final states, may be related to the quantum ergodic theorem. Systems possessing the quantum ergodicity property as defined in [11] are those systems "where all states, within a given energy range can be reached from all other states within the range even if only via other intermediate states". In our case, if quantum ergodicity holds, it will be possible to travel from any $|\psi_i\rangle$ to any $|\psi_f\rangle$ via two different routes.

Substituting Schrodinger's equation directly into EL equation would have been the easier way to show the validity of the principle. However, we highlight equation 7 to point out the intriguing possibility that some nonlinear equations may satisfy this equation and consequently fulfills Fermat principle. This is indeed the case, and we found that the nonlinear equation

$$|\dot{\psi}\rangle = \pm iH|\psi\rangle \mp 2i\,\langle\psi|H|\psi\rangle\,|\psi\rangle \tag{8}$$

satisfies equation 7.

As emphasized above, the timeless nature of this variational principle does not make it adequate to deal with time-dependent Hamiltonians. To achieve this aim, we need to add a 'time-like' component to the Fubini-Study metric and modify the velocity term $\nu(s)$ accordingly. There exists a variety of ways to implement this prescription. We adopt the following choice of the full metric, $ds'^2 = ds^2 - \langle\psi|H(t)^2|\psi\rangle\,dt^2$ and the speed term $\nu(s)$



FIG. 1: **Generalized Fermat Principle**. Natural evolution of a system from initial to final state (thick) in the projective Hilbert space (a), on the energy shell in classical phase space (b) and in the classical configuration space (c) together with fictitious paths (dashed) connecting the same states. The time functional $\mathcal{T}$ is an extremum along the physical path in case (a) and (b) but in general not in (c). The path is a geodesic on the energy shell in (b) but not in (a) and (c).

becomes $\langle\psi|H(t)|\psi\rangle$. After substituting into equation 6 with the new integrand and taking the inner product with $|\psi\rangle$, we arrive at

$$\langle\dot{\psi}|\dot{\psi}\rangle = \langle\dot{\psi}|\psi\rangle^2 + |\langle\dot{\psi}|\psi\rangle|^2 + \langle\psi|H(t)^2|\psi\rangle \tag{9}$$

We notice that both Schrodinger's equation and the nonlinear equation 8 satisfy this equation. The extension of the FS metric to the new one is analogous to the transition from the Maupertuis action $\int pdq$ to the Hamiltonian action in classical mechanics $\int dt\,(p\dot{q} - H(p,q,t))$. Had we chosen the new metric to be $ds'^2 = ds^2 + \langle\psi|H(t)|\psi\rangle^2\, dt^2$ the new velocity term would have become $\langle\psi|H(t)^2|\psi\rangle^{\frac{1}{2}}$. These time dependent variational principles are to be contrasted with the functional $\langle\psi(t)|H(t) - i\partial_t|\psi(t)\rangle$ from which Schrodinger's equation can be derived. The later however lacks an appealing geometric interpretation.

—Fermat Principle in Phase Space We consider a conservative classical system composed of N particles described by the generalized coordinates q_i, p_i, with velocity-independent potential $V(q_1, ...q_N)$, kinetic energy $T = \sum_{i=1}^{N} p_i^2$ and Hamiltonian $H = T + V$. Let the distance element in phase space be described by the Euclidean metric $ds^2 = \sum_{i=1}^{N} dq_i^2 + dp_i^2$. The speed along the physical path generated by Hamiltonian flow, $\nu(s)$ equals $\sqrt{\sum_{i=1}^{N}\left(\frac{\partial H}{\partial q_i}\right)^2 + \left(\frac{\partial H}{\partial p_i}\right)^2}$ [11]. We therefore express the time functional $\mathcal{T}$ as

$$\mathcal{T} = \int_{x_i}^{x_f} \frac{ds}{\sqrt{\sum_i \left(\frac{\partial H}{\partial x_i}\right)^2}} \tag{10}$$

where x_i denotes all generalized coordinates q_i, p_i treated on equal footing. Adding quantities of different dimensions in the metric and velocity terms should not disturb the reader because it simply reflects the fact that the axes of the phase space $\{q_i, p_i\}$ have different dimensions. Depending on the particular problem, any of $\{q_i, p_i\}$ can be redefined in terms of the physical constants of the problem so that they have the same dimensions.

We investigate Fermat principle in phase space on the energy shell $H(x_i) = E$ (see Fig. 1.b), in contrast to the quantum case where we didn't have to impose that constraint. That translates mathematically into the variational problem $\delta\mathcal{T} = 0$ subject to the constraint $H(x_i) - E = 0$. If we parametrize any arbitrary trajectory connecting x_i and x_f by τ, we can define the integrand L which satisfies the EL equation

$$\frac{\delta L}{\delta x} - \frac{d}{d\tau}\frac{\delta L}{\delta \dot{x}} = 0 \tag{11}$$

to be $L = \lambda(\tau)H'(x_i) + F(x_i)G(\dot{x_i})$, where $H' = H - E$, $F = \frac{1}{\sqrt{\sum_i \left(\frac{\partial H}{\partial x_i}\right)^2}}$ and $G = \sqrt{\sum_i \dot{x_i}^2}$. Substituting into equation 11, we end up with

$$\lambda(\tau)\frac{\partial H'}{\partial x_i}G + G^2\frac{\partial F}{\partial x_i} - \frac{G^2\left(\ddot{x}_i F + \dot{x}_i \sum_j \frac{\partial F}{\partial x_j}\dot{x_j}\right) - \dot{x}_i F \sum_j \dot{x}_j \ddot{x}_j}{G^2} = 0. \tag{12}$$

If Fermat principle holds in phase space, the previous equation has to be satisfied for every x_i. By multiplying each equation by $\dot{x_i}$ and adding up all of them, we end up with

$$\lambda(\tau)G\sum_i \dot{x_i}\frac{\partial H'}{\partial x_i} = 0. \tag{13}$$

We see immediately that the Hamiltonian equations of motion $\dot{q_i} = \frac{\partial H}{\partial p_i}$ and $\dot{p_i} = -\frac{\partial H}{\partial q_i}$ satisfy this equation, when τ is taken to be the real time. As in the quantum case, the time-reversed evolution $\dot{q_i} = -\frac{\partial H}{\partial p_i}$ and $\dot{p_i} = \frac{\partial H}{\partial q_i}$ will also satisfy this equation. For non-periodic systems, that would be of relevance to the classical ergodicity problem, where the system starting from any point in phase space will pass arbitrarily close to any other point on the same energy shell via forward and backward time evolution. It is easy to check after solving the Hamilton's equations of motion for a given system and substituting back in equation 12 that we get consistent solutions for $\lambda(\tau)$ from all the equations.

We point out that by the same procedure one can show that the phase space path length $\int ds$ is also a stationary quantity under the constraint of conserved energy for Hamiltonian flow, i.e., unlike configuration space, the system follows a geodesic path, at least locally, on the enrgy shell in phase space. One interesting question to ask is whether the phase space path can be determined uniquely by extremizing $\int ds$ under time constraint rather than energy constraint. The time constraint is $\int \frac{ds}{\nu(s)} = T$ and it is an isoperimetric constraint rather than a holonomic constraint like the former case. The answer is No! One can check for any simple system that after substituting in the EL equation with the solution for $x_i(\tau)$ generated by the Hamiltonian flow, that one does not get consistent solution for the Lagrange multiplier $\lambda(\tau)$ for every x_i. Therefore, counterintuitively, time constraint is not equivalent to energy constraint in phase space.

In classical systems where kinetic and potential energies cannot be straightforwardly defined, e.g., classical spin systems, we hypothesize that Fermat principle will continue to hold in the proper space that completely specify the state of the system. In case of N interacting classical spins, the phase space would be a 2N-dimensional manifold where each spin is specified by two angles (θ_i, ϕ_i) and the conjugate variables $\{q_i, p_i\}$ would correspond to $\{cos(\theta_i), \phi_i\}$ [12].

We finally comment on the validity of Fermat Principle in the configuration space of a conservative classical many-body system described by the coordinates $\{q_i\}$ (Fig. 1.c). The metric is defined by the kinetic energy term which we assume to be quadratic in velocities, i.e., $T = \frac{1}{2}m_{ij}q_iq_j$. It is easy to verify that T is related to the speed $\nu(s)$ of the Hamiltonian flow in the configuration space by $\nu(s) = \sqrt{2T}$. The validity of Fermat Principle in configuration space would imply that the action $\int \frac{ds}{\sqrt{T}}$ is equivalent to the Jacobi action $\int \sqrt{T}ds$, i.e., yields the same equations of motion. To test this supposition, we compare the Euler-Lagrange equations that will result from extremizing the two actions. Let $ds = F(q, \dot{q})\,d\tau$ and $\sqrt{T} = G(q, \dot{q}, \tau)$. We can see by straightforward substitution into $\frac{\delta L}{\delta q_i} - \frac{d}{dt}\frac{\delta L}{\delta \dot{q_i}} = 0$ where L is the integrand of the action, that the two actions $\int \frac{F}{G}d\tau$ and $\int FGd\tau$ will satisfy the same equation of motion if $F(q, \dot{q})$ satisfy $\frac{\delta F}{\delta q_i} - \frac{d}{dt}\frac{\delta F}{\delta \dot{q_i}} = 0$. That means that $\int ds$ should itself be a stationary quantity along the Hamiltonian flow, or in other words the system follows a geodesic in the bare configuration space. Since the flow is not geodesic in this space except for the case of zero potential energy [4] (or more generally when the system is confined to an equi-potential energy surface in the configuration space), we conclude that Fermat principle is generally not valid in this case for a non-vanishing potential , i.e., $\delta\mathcal{T} \neq 0$[13]. To make the previous argument more rigorous, one should take into consideration while solving the EL equation and comparing the two functionals the constraint that the potential energy is always smaller than the total energy, $V(q_1, q_2, ...) < E$, but we suppose that the same conclusion will be reached.

In conclusion, we have introduced a generalized Fermat principle as a unified concept in Hilbert and phase spaces of quantum and classical systems that illuminates new aspects of Fermat's surmise of *natural economy*. The generalized Fermat principle provides new geometric variational principles that are satisfied naturally by the Schrodinger's and Hamilton's equations of motion.

The concepts developed in this study provokes several questions which are left open. What are the implications of the generalized Fermat principle on the nature of time in classical and quantum dynamics? Why is it natural to expect that the equations of dynamical evolution in a complete space should yield longer times along all paths other than the actual path? Can this observation be of any practical use to simulate complex many body systems that require unaffordable computing resources, or at least get insight into their behavior? Can we use the quantum Fermat principle to define a new basis where the quantum evolution follows geodesic paths? What are its implications for quantum computing? Are there situations where the nonlinear equation of motion for the wave function (8) that satisfies Fermat principle be physically relevant? What does Fermat principle say about the statistical evolution of an ensemble of systems in phase space? Can it be used to regard Liouville's equation as a variational principle? We hope these questions will motivate more researchers in the future to investigate the nature of the geometry of evolution in classical and quantum mechanics.

The author would like to thank Boris Fine for many fruitful discussions throughout the development of this paper. The comments of Chris Gray are highly appreciated.

---

[*] Electronic address: T.Elsayed@thphys.uni-heidelberg.de

[1] C. G. Gray, G. Karl, and V. A. Novikov, Reports on Progress in Physics **67**, 159 (2004).
[2] J.-L. Basdevant, *Variational Principles in Physics* (Springer, 2007).
[3] W. Yourgrau and S. Mandelstam, *Variational Principles in Dynamics and Quantum Theory* (Courier Dover Publications, 1979).
[4] C. Lanczos, *The Variational Principles of Mechanics* (Courier Dover Publications, 1970).
[5] Y. Aharonov and J. Anandan, Phys. Rev. Lett. **58**, 1593 (1987).
[6] J. Anandan and Y. Aharonov, Phys. Rev. Lett. **65**, 1697 (1990).
[7] J. Anandan and Y. Aharonov, Phys. Rev. D **38**, 1863 (1988).
[8] A. Kryukov (2007), published in Foundations of Physics, 37, (2007), 3.
[9] A. N. Grigorenko, Phys. Rev. A **46**, 7292 (1992).
[10] A. Carlini, A. Hosoya, T. Koike, and Y. Okudaira, Phys. Rev. Lett. **96**, 060503 (2006).
[11] D. T. Haar, Rev. Mod. Phys. **27**, 289 (1955).
[12] R. Steinigeweg and H.-J. Schmidt, Computer Physics Communications **174**, 853 (2006).
[13] S. Gryb, Phys. Rev. D **81**, 044035 (2010).